

Branching Ratio and CP Asymmetry of $B \rightarrow \pi\eta^{(\prime)}$ Decays in the Perturbative QCD Approach

Huisheng Wang, Xin Liu, Zhenjun Xiao (corresponding author),* and Libo Guo†

*Department of Physics and Institute of Theoretical Physics,
Nanjing Normal University, Nanjing, Jiangsu 210097, P.R.China*

Cai-Dian Lü‡

*CCAST(World Laboratory), P.O. Box 8730,
Beijing 100080, P.R. China; and*

*Institute of High Energy Physics, CAS,
P.O.Box 918(4) Beijing 100049, China §*

(Dated: February 2, 2008)

Abstract

We calculate the branching ratios and CP-violating asymmetries for $B^0 \rightarrow \pi^0\eta^{(\prime)}$ and $B^+ \rightarrow \pi^+\eta^{(\prime)}$ decays in the perturbative QCD (pQCD) factorization approach here. We not only calculate the usual factorizable contributions, but also evaluate the non-factorizable and annihilation type contributions. Besides the current-current operators, the contributions from the QCD and electroweak penguin operators are also taken into account. The pQCD results for the CP-averaged branching ratios are $Br(B^+ \rightarrow \pi^+\eta) \approx 4.1 \times 10^{-6}$, $Br(B^+ \rightarrow \pi^+\eta') \approx 2.4 \times 10^{-6}$, and $Br(B^0 \rightarrow \pi^0\eta^{(\prime)}) \approx 0.2 \times 10^{-6}$, which agree very well with the measured values or currently available experimental upper limits. We also predict large CP-violating asymmetries in these decays: $A_{CP}^{dir}(\pi^\pm\eta^{(\prime)}) \sim A_{CP}^{dir}(\pi^0\eta^{(\prime)}) \sim -0.35$, and $A_{CP}^{mix}(\pi^0\eta^{(\prime)}) \sim 0.67$, but with large errors. The pQCD prediction for $A_{CP}^{dir}(\pi^\pm\eta)$ ($A_{CP}^{dir}(\pi^\pm\eta')$) has the same (opposite) sign with the primary measured values. Further improvements in both theory and experiments are needed to clarify this discrepancy.

PACS numbers: 13.25.Hw, 12.38.Bx, 14.40.Nd

§ Mailing address.

*Electronic address: xiaozhenjun@njnu.edu.cn

†Electronic address: guolb@email.njnu.edu.cn

‡Electronic address: lucd@ihep.ac.cn

I. INTRODUCTION

The experimental measurements and theoretical studies of the two body charmless hadronic B meson decays play an important role in the precision test of the standard mode (SM) and in searching for the new physics beyond the SM [1]. For these charmless B meson decays, the dominant theoretical error comes from the large uncertainty in evaluating the hadronic matrix elements $\langle M_1 M_2 | O_i | B \rangle$ where M_1 and M_2 are light final state mesons. The QCD factorization (QCDF) approach [2] and the perturbative QCD (pQCD) factorization approach [3, 4, 5] are the popular methods being used to calculate the hadronic matrix element. Many charmless B meson decays, for example, have been calculated in the QCDF approach [2, 6, 7, 8] and/or in the pQCD approach [9, 10, 11, 12, 13, 14].

In this paper, we would like to calculate the branching ratios and CP asymmetries for the four $B \rightarrow \pi \eta^{(\prime)}$ decays by employing the low energy effective Hamiltonian [15] and the pQCD approach. Besides the usual factorizable contributions, we here are able to evaluate the non-factorizable and the annihilation contributions to these decays. Theoretically, the four $B \rightarrow \pi \eta^{(\prime)}$ decays have been studied before in the naive or generalized factorization approach [16] as well as in the QCDF approach [6]. On the experimental side, the CP-averaged branching ratios and CP-violating asymmetries of $B \rightarrow \pi^+ \eta$ and $\pi^+ \eta'$ decays have been measured very recently [17, 18], and the world averages as given by the Heavy Flavor Averaging Group [19] are

$$Br(B^+ \rightarrow \pi^+ \eta) = (4.3 \pm 0.4) \times 10^{-6}, \quad (1)$$

$$Br(B^+ \rightarrow \pi^+ \eta') = (2.53^{+0.59}_{-0.50}) \times 10^{-6}, \quad (2)$$

$$\mathcal{A}_{CP}(B^\pm \rightarrow \pi^\pm \eta) = -0.11 \pm 0.08, \quad (3)$$

$$\mathcal{A}_{CP}(B^+ \rightarrow \pi^+ \eta') = 0.14 \pm 0.15. \quad (4)$$

For $B \rightarrow \pi^0 \eta, \pi^0 \eta'$ decays, only the experimental upper limits for the branching ratios are available now [19]

$$Br(B^0 \rightarrow \pi^0 \eta) < 2.5 \times 10^{-6}, \quad Br(B^0 \rightarrow \pi^0 \eta') < 3.7 \times 10^{-6}. \quad (5)$$

For $B \rightarrow \pi \eta^{(\prime)}$ decays, the B meson is heavy, setting at rest and decaying into two light mesons (i.e. π and $\eta^{(\prime)}$) with large momenta. Therefore the light final state mesons are moving very fast in the rest frame of B meson. In this case, the short distance hard process dominates the decay amplitude. We shall demonstrate that the soft final state interaction is not important for such decays, since there is not enough time for light mesons to exchange soft gluons. Therefore, it makes the pQCD reliable in calculating the $B \rightarrow \pi \eta^{(\prime)}$ decays. With the Sudakov resummation, we can include the leading double logarithms for all loop diagrams, in association with the soft contribution. Unlike the usual factorization approach, the hard part of the pQCD approach consists of six quarks rather than four. We thus call it six-quark operators or six-quark effective theory. Applying the six-quark effective theory to B meson decays, we need meson wave functions for the hadronization of quarks into mesons. All the collinear dynamics are included in the meson wave functions.

This paper is organized as follows. In Sec. II, we give a brief review for the pQCD factorization approach. In Sec. III, we calculate analytically the related Feynman diagrams and present the various decay amplitudes for the studied decay modes. In Sec. IV, we

show the numerical results for the branching ratios and CP asymmetries of $B \rightarrow \pi\eta^{(\prime)}$ decays and compare them with the measured values or the theoretical predictions in QCDF approach. The summary and some discussions are included in the final section.

II. THEORETICAL FRAMEWORK

The three scale PQCD factorization approach has been developed and applied in the non-leptonic B meson decays [3, 4, 5, 9, 10, 11, 12] for some time. In this approach, the decay amplitude is separated into soft, hard, and harder dynamics characterized by different energy scales (t, m_b, M_W). It is conceptually written as the convolution,

$$\mathcal{A}(B \rightarrow M_1 M_2) \sim \int d^4 k_1 d^4 k_2 d^4 k_3 \text{Tr} [C(t) \Phi_B(k_1) \Phi_{M_1}(k_2) \Phi_{M_2}(k_3) H(k_1, k_2, k_3, t)], \quad (6)$$

where k_i 's are momenta of light quarks included in each mesons, and Tr denotes the trace over Dirac and color indices. $C(t)$ is the Wilson coefficient which results from the radiative corrections at short distance. In the above convolution, $C(t)$ includes the harder dynamics at larger scale than M_B scale and describes the evolution of local 4-Fermi operators from m_W (the W boson mass) down to $t \sim \mathcal{O}(\sqrt{\bar{\Lambda} M_B})$ scale, where $\bar{\Lambda} \equiv M_B - m_b$. The function $H(k_1, k_2, k_3, t)$ describes the four quark operator and the spectator quark connected by a hard gluon whose q^2 is in the order of $\bar{\Lambda} M_B$, and includes the $\mathcal{O}(\sqrt{\bar{\Lambda} M_B})$ hard dynamics. Therefore, this hard part H can be perturbatively calculated. The function Φ_M is the wave function which describes hadronization of the quark and anti-quark to the meson M . While the function H depends on the processes considered, the wave function Φ_M is independent of the specific processes. Using the wave functions determined from other well measured processes, one can make quantitative predictions here.

Since the b quark is rather heavy we consider the B meson at rest for simplicity. It is convenient to use light-cone coordinate $(p^+, p^-, \mathbf{p}_T)$ to describe the meson's momenta,

$$p^\pm = \frac{1}{\sqrt{2}}(p^0 \pm p^3), \quad \text{and} \quad \mathbf{p}_T = (p^1, p^2). \quad (7)$$

Using these coordinates the B meson and the two final state meson momenta can be written as

$$P_1 = \frac{M_B}{\sqrt{2}}(1, 1, \mathbf{0}_T), \quad P_2 = \frac{M_B}{\sqrt{2}}(1, 0, \mathbf{0}_T), \quad P_3 = \frac{M_B}{\sqrt{2}}(0, 1, \mathbf{0}_T), \quad (8)$$

respectively, here the light meson masses have been neglected. Putting the light (anti-) quark momenta in B , π and η mesons as k_1 , k_2 , and k_3 , respectively, we can choose

$$k_1 = (x_1 P_1^+, 0, \mathbf{k}_{1T}), \quad k_2 = (x_2 P_2^+, 0, \mathbf{k}_{2T}), \quad k_3 = (0, x_3 P_3^-, \mathbf{k}_{3T}). \quad (9)$$

Then, the integration over k_1^- , k_2^- , and k_3^+ in eq.(6) will lead to

$$\begin{aligned} \mathcal{A}(B \rightarrow \pi\eta^{(\prime)}) \sim & \int dx_1 dx_2 dx_3 b_1 db_1 b_2 db_2 b_3 db_3 \\ & \cdot \text{Tr} [C(t) \Phi_B(x_1, b_1) \Phi_\pi(x_2, b_2) \Phi_{\eta^{(\prime)}}(x_3, b_3) H(x_i, b_i, t) S_t(x_i) e^{-S(t)}] \end{aligned} \quad (10)$$

where b_i is the conjugate space coordinate of k_{iT} , and t is the largest energy scale in function $H(x_i, b_i, t)$. The large logarithms ($\ln m_W/t$) coming from QCD radiative corrections to four quark operators are included in the Wilson coefficients $C(t)$. The large double logarithms ($\ln^2 x_i$) on the longitudinal direction are summed by the threshold resummation [20], and they lead to $S_t(x_i)$ which smears the end-point singularities on x_i . The last term, $e^{-S(t)}$, is the Sudakov form factor which suppresses the soft dynamics effectively [21]. Thus it makes the perturbative calculation of the hard part H applicable at intermediate scale, i.e., M_B scale. We will calculate analytically the function $H(x_i, b_i, t)$ for $B \rightarrow \pi\eta^{(\prime)}$ decays in the first order in α_s expansion and give the convoluted amplitudes in next section.

A. Wilson Coefficients

For $B \rightarrow \pi\eta^{(\prime)}$ decays, the related weak effective Hamiltonian H_{eff} can be written as [15]

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \left[V_{ub}V_{ud}^* (C_1(\mu)O_1^u(\mu) + C_2(\mu)O_2^u(\mu)) - V_{tb}V_{td}^* \sum_{i=3}^{10} C_i(\mu) O_i(\mu) \right]. \quad (11)$$

We specify below the operators in $\mathcal{H}_{eff}$ for $b \rightarrow d$ transition:

$$\begin{aligned} O_1^u &= \bar{d}_\alpha \gamma^\mu L u_\beta \cdot \bar{u}_\beta \gamma_\mu L b_\alpha, & O_2^u &= \bar{d}_\alpha \gamma^\mu L u_\alpha \cdot \bar{u}_\beta \gamma_\mu L b_\beta, \\ O_3 &= \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu L q'_\beta, & O_4 &= \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu L q'_\alpha, \\ O_5 &= \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu R q'_\beta, & O_6 &= \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu R q'_\alpha, \\ O_7 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu R q'_\beta, & O_8 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu R q'_\alpha, \\ O_9 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu L q'_\beta, & O_{10} &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu L q'_\alpha, \end{aligned} \quad (12)$$

where α and β are the $SU(3)$ color indices; L and R are the left- and right-handed projection operators with $L = (1 - \gamma_5)$, $R = (1 + \gamma_5)$. The sum over q' runs over the quark fields that are active at the scale $\mu = O(m_b)$, i.e., ($q' \in \{u, d, s, c, b\}$). The PQCD approach works well for the leading twist approximation and leading double logarithm summation. For the Wilson coefficients $C_i(\mu)$ ($i = 1, \dots, 10$), we will also use the leading order (LO) expressions, although the next-to-leading order (NLO) calculations already exist in the literature [15]. This is the consistent way to cancel the explicit μ dependence in the theoretical formulae.

For the renormalization group evolution of the Wilson coefficients from higher scale to lower scale, we use the formulae as given in Ref.[9] directly. At the high m_W scale, the leading order Wilson coefficients $C_i(M_W)$ are simple and can be found easily in Ref.[15]. In PQCD approach, the scale t may be larger or smaller than the m_b scale. For the case of $m_b < t < m_W$, we evaluate the Wilson coefficients at t scale using leading logarithm running equations, as given in Eq.(C1) of Ref.[9]. In numerical calculations, we use $\alpha_s = 4\pi/[\beta_1 \ln(t^2/\Lambda_{QCD}^{(5)})]$ which is the leading order expression with $\Lambda_{QCD}^{(5)} = 193\text{MeV}$, derived from $\Lambda_{QCD}^{(4)} = 250\text{MeV}$. Here $\beta_1 = (33 - 2n_f)/12$, with the appropriate number of active quarks n_f . $n_f = 5$ when scale t is larger than m_b .

At a given energy scale $t = m_b = 4.8$ GeV, the LO Wilson coefficients $C_i(m_b)$ as given in Ref.[9] are

$$\begin{aligned} C_1 &= -0.2703, & C_2 &= 1.1188, & C_3 &= 0.0126, & C_4 &= -0.0270, \\ C_5 &= 0.0085, & C_6 &= -0.0326, & C_7 &= 0.0011, & C_8 &= 0.0004, \\ C_9 &= -0.0090, & C_{10} &= 0.0022. \end{aligned} \quad (13)$$

If the scale $t < m_b$, then we evaluate the Wilson coefficients at t scale using the input of Eq. (13) and the formulae in Appendix D of Ref.[9] for the case of $n_f = 4$.

B. Wave Functions

In the resummation procedures, the B meson is treated as a heavy-light system. In general, the B meson light-cone matrix element can be decomposed as [22, 23]

$$\begin{aligned} & \int_0^1 \frac{d^4 z}{(2\pi)^4} e^{i\mathbf{k}_1 \cdot \mathbf{z}} \langle 0 | \bar{b}_\alpha(0) d_\beta(z) | B(p_B) \rangle \\ &= -\frac{i}{\sqrt{2N_c}} \left\{ (\not{p}_B + m_B) \gamma_5 \left[\phi_B(\mathbf{k}_1) - \frac{\not{n} - \not{v}}{\sqrt{2}} \bar{\phi}_B(\mathbf{k}_1) \right] \right\}_{\beta\alpha}, \end{aligned} \quad (14)$$

where $n = (1, 0, \mathbf{0}_T)$, and $v = (0, 1, \mathbf{0}_T)$ are the unit vectors pointing to the plus and minus directions, respectively. From the above equation, one can see that there are two Lorentz structures in the B meson distribution amplitudes. They obey to the following normalization conditions

$$\int \frac{d^4 k_1}{(2\pi)^4} \phi_B(\mathbf{k}_1) = \frac{f_B}{2\sqrt{2N_c}}, \quad \int \frac{d^4 k_1}{(2\pi)^4} \bar{\phi}_B(\mathbf{k}_1) = 0. \quad (15)$$

In general, one should consider these two Lorentz structures in calculations of B meson decays. However, it can be argued that the contribution of $\bar{\phi}_B$ is numerically small [24, 25], thus its contribution can be numerically neglected. In this approximation, we keep minimum number of input parameters for wave functions. Therefore, we only consider the contribution of Lorentz structure

$$\Phi_B = \frac{1}{\sqrt{2N_c}} (\not{p}_B + m_B) \gamma_5 \phi_B(\mathbf{k}_1), \quad (16)$$

in our calculation. We use the same wave functions as in Refs.[9, 10, 25, 26]. In the next section, we will see that the hard part is always independent of one of the k_1^+ and/or k_1^- , if we make some approximations. The B meson wave function is then the function of variable k_1^- (or k_1^+) and $k_1^\perp$.

$$\phi_B(k_1^-, k_1^\perp) = \int dk_1^+ \phi(k_1^+, k_1^-, k_1^\perp). \quad (17)$$

The wave function for $d\bar{d}$ components in π meson are given as

$$\Phi_\pi(P, x, \zeta) \equiv \frac{1}{\sqrt{2N_C}} \left[\not{p} \phi_\pi(x) + m_0^\pi \phi_\pi^P(x) + \zeta m_0^\pi (\not{p} \not{v} - v \cdot n) \phi_\pi^T(x) \right]. \quad (18)$$

The wave function for $d\bar{d}$ components of $\eta^{(\prime)}$ meson are given as

$$\Phi_{\eta_{d\bar{d}}}(P, x, \zeta) \equiv \frac{1}{\sqrt{2N_C}} \left[\not{p} \phi_{\eta_{d\bar{d}}}^A(x) + m_0^{\eta_{d\bar{d}}} \phi_{\eta_{d\bar{d}}}^P(x) + \zeta m_0^{\eta_{d\bar{d}}} (\not{p} \not{\eta} - v \cdot n) \phi_{\eta_{d\bar{d}}}^T(x) \right], \quad (19)$$

where P and x are the momentum and the momentum fraction of $\eta_{d\bar{d}}$, respectively. We assumed here that the wave function of $\eta_{d\bar{d}}$ is same as the π wave function. The parameter ζ is either $+1$ or -1 depending on the assignment of the momentum fraction x . The $s\bar{s}$ component of the wave function can be similarly defined.

The transverse momentum $k^\perp$ is usually conveniently converted to the b parameter by Fourier transformation. The initial conditions of leading twist $\phi_i(x)$, $i = B, \pi, \eta, \eta'$, are of non-perturbative origin, satisfying the normalization

$$\int_0^1 \phi_i(x, b=0) dx = \frac{1}{2\sqrt{6}} f_i, \quad (20)$$

with f_i the meson decay constants.

III. PERTURBATIVE CALCULATIONS

In the previous section we have discussed the wave functions and Wilson coefficients of the amplitude in eq.(6). In this section, we will calculate the hard part $H(t)$. This part involves the four quark operators and the necessary hard gluon connecting the four quark operator and the spectator quark. We will show the whole amplitude for each diagram including wave functions. Similar to the $B \rightarrow \pi\rho$ [27] and $B \rightarrow \rho\eta^{(\prime)}$ decays [14], there are 8 type diagrams contributing to the $B \rightarrow \pi\eta^{(\prime)}$ decays, as illustrated in Figure 1. We first calculate the usual factorizable diagrams (a) and (b). Operators O_1, O_2, O_3, O_4, O_9 , and O_{10} are $(V-A)(V-A)$ currents, the sum of their amplitudes is given as

$$\begin{aligned} F_{e\pi} = & 8\pi C_F m_B^4 \int_0^1 dx_1 dx_3 \int_0^\infty db_1 db_3 db_3 \phi_B(x_1, b_1) \\ & \times \left\{ \left[(1+x_3) \phi_\pi(x_3, b_3) + (1-2x_3) r_\pi (\phi_\pi^p(x_3, b_3) + \phi_\pi^t(x_3, b_3)) \right] \right. \\ & \cdot \alpha_s(t_e^1) h_e(x_1, x_3, b_1, b_3) \exp[-S_{ab}(t_e^1)] \\ & \left. + 2r_\pi \phi_\pi^p(x_3, b_3) \alpha_s(t_e^2) h_e(x_3, x_1, b_3, b_1) \exp[-S_{ab}(t_e^2)] \right\}, \end{aligned} \quad (21)$$

where $r_\pi = m_0^\pi/m_B$; $C_F = 4/3$ is a color factor. The function h_e , the scales t_e^i and the Sudakov factors S_{ab} are displayed in Appendix A. In the above equation, we do not include the Wilson coefficients of the corresponding operators, which are process dependent. They will be shown later in this section for different decay channels.

The form factors of B to π decay, $F_{0,1}^{B \rightarrow \pi}(0)$, can thus be extracted from the expression in Eq. (21), that is

$$F_0^{B \rightarrow \pi}(q^2=0) = F_1^{B \rightarrow \pi}(q^2=0) = F_{e\pi}/m_B^2, \quad (22)$$

which is identical with that defined in Ref.[24].

The operators O_5, O_6, O_7 , and O_8 have a structure of $(V-A)(V+A)$. In some decay channels, some of these operators contribute to the decay amplitude in a factorizable way.




FIG. 1: Diagrams contributing to the $B \rightarrow \pi \eta^{(\prime)}$ decays (diagram (a) and (b) contribute to the $B \rightarrow \pi$ form factor $F_{0,1}^{B \rightarrow \pi}$).

Since only the axial-vector part of $(V + A)$ current contribute to the pseudo-scalar meson production, $\langle \pi | V - A | B \rangle \langle \eta | V + A | 0 \rangle = -\langle \pi | V - A | B \rangle \langle \eta | V - A | 0 \rangle$, that is

$$F_{e\pi}^{P1} = -F_{e\pi} . \quad (23)$$

In some other cases, we need to do Fierz transformation for these operators to get right color structure for factorization to work. In this case, we get $(S - P)(S + P)$ operators from $(V - A)(V + A)$ ones. For these $(S - P)(S + P)$ operators, Fig. 1(a) and 1(b) give

$$\begin{aligned} F_{e\pi}^{P2} = & -16\pi C_F m_B^4 r_\pi \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) \\ & \times \left\{ \left[\phi_\pi(x_3, b_3) + r_\pi((2 + x_3)\phi_\pi^P(x_3, b_3) - x_3\phi_\pi^t(x_3, b_3)) \right] \right. \\ & \cdot \alpha_s(t_e^1) h_e(x_1, x_3, b_1, b_3) \exp[-S_{ab}(t_e^1)] \\ & + \left[x_1\phi_\pi(x_3, b_3) - 2(x_1 - 1)r_\pi\phi_\pi^P(x_3, b_3) \right] \\ & \left. \cdot \alpha_s(t_e^2) h_e(x_3, x_1, b_3, b_1) \exp[-S_{ab}(t_e^2)] \right\} . \end{aligned} \quad (24)$$

For the non-factorizable diagrams 1(c) and 1(d), all three meson wave functions are involved. The integration of b_3 can be performed using δ function $\delta(b_3 - b_2)$, leaving only integration of b_1 and b_2 . For the $(V - A)(V - A)$ operators, the result is

$$M_{e\pi} = \frac{16\sqrt{6}}{3}\pi C_F m_B^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \phi_\eta(x_2, b_2) \\ \times x_3 [2r_\pi \phi_\pi^t(x_3, b_2) - \phi_\pi(x_3, b_2)] \\ \cdot \alpha_s(t_f) h_f(x_1, x_2, x_3, b_1, b_2) \exp[-S_{cd}(t_f)] . \quad (25)$$

For the $(V - A)(V + A)$ operators the formulae are different. Here we have two kinds of contributions from $(V - A)(V + A)$ operators. $M_{e\pi}^{P1}$ and $M_{e\pi}^{P2}$ is for the $(V - A)(V + A)$ and $(S - P)(S + P)$ type operators respectively:

$$M_{e\pi}^{P1} = 0, \quad (26)$$

$$M_{e\pi}^{P2} = -M_{e\pi} . \quad (27)$$

The factorizable annihilation diagrams (g) and (h) involve only π and $\eta^{(\prime)}$ wave functions. There are also three kinds of decay amplitudes for these two diagrams. $F_{a\pi}$ is for $(V - A)(V - A)$ type operators, $F_{a\pi}^{P1}$ is for $(V - A)(V + A)$ type operators, while $F_{a\pi}^{P2}$ is for $(S - P)(S + P)$ type operators:

$$F_{a\pi}^{P1} = F_{a\pi} = 8\pi C_F m_B^4 \int_0^1 dx_2 dx_3 \int_0^\infty b_2 db_2 b_3 db_3 \{ [x_3 \phi_\pi(x_3, b_3) \phi_\eta(x_2, b_2) \\ + 2r_\eta r_\pi ((x_3 + 1) \phi_\pi^P(x_3, b_3) + (x_3 - 1) \phi_\pi^t(x_3, b_3)) \phi_\eta^P(x_2, b_2)] \\ \cdot \alpha_s(t_e^3) h_a(x_2, x_3, b_2, b_3) \exp[-S_{gh}(t_e^3)] \\ - [x_2 \phi_\pi(x_3, b_3) \phi_\eta(x_2, b_2) \\ + 2r_\eta r_\pi \phi_\pi^P(x_3, b_3) ((x_2 + 1) \phi_\eta^P(x_2, b_2) + (x_2 - 1) \phi_\eta^t(x_2, b_2))] \\ \cdot \alpha_s(t_e^4) h_a(x_3, x_2, b_3, b_2) \exp[-S_{gh}(t_e^4)] \} , \quad (28)$$

$$F_{a\pi}^{P2} = 16\pi C_F m_B^4 \int_0^1 dx_2 dx_3 \int_0^\infty b_2 db_2 b_3 db_3 \\ \times \{ [x_3 r_\pi (\phi_\pi^P(x_3, b_3) - \phi_\pi^t(x_3, b_3)) \phi_\eta(x_2, b_2) + 2r_\eta \phi_\pi(x_3, b_3) \phi_\eta^P(x_2, b_2)] \\ \times \alpha_s(t_e^3) h_a(x_2, x_3, b_2, b_3) \exp[-S_{gh}(t_e^3)] \\ + [2r_\pi \phi_\pi^P(x_3, b_3) \phi_\eta(x_2, b_2) + x_2 r_\eta (\phi_\eta^P(x_2, b_2) - \phi_\eta^t(x_2, b_2)) \phi_\pi(x_3, b_3)] \\ \times \alpha_s(t_e^4) h_a(x_3, x_2, b_3, b_2) \exp[-S_{gh}(t_e^4)] \} . \quad (29)$$

For the non-factorizable annihilation diagrams (e) and (f), again all three wave functions are involved. Here we have two kinds of contributions. $M_{a\pi}$, $M_{a\pi}^{P1}$ and $M_{a\pi}^{P2}$ describe the contributions from the $(V - A)(V - A)$, $(V - A)(V + A)$ and $(S - P)(S + P)$ type

operators, respectively,

$$\begin{aligned}
M_{a\pi} = & \frac{16\sqrt{6}}{3}\pi C_F m_B^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \\
& \times \{ -\{x_2 \phi_\pi(x_3, b_2) \phi_\eta(x_2, b_2) \\
& + r_\pi r_\eta [(x_2 + x_3 + 2) \phi_\eta^P(x_2, b_2) + (x_2 - x_3) \phi_\eta^t(x_2, b_2)] \phi_\pi^P(x_3, b_2) \\
& + [(x_2 - x_3) \phi_\pi^P(x_3, b_2) + (x_2 + x_3 - 2) \phi_\eta^t(x_2, b_2)] \phi_\pi^t(x_3, b_2) \} \\
& \cdot \alpha_s(t_f^3) h_f^3(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^3)] \\
& + \{x_3 \phi_\pi(x_3, b_2) \phi_\eta(x_2, b_2) \\
& - r_\pi r_\eta [\phi_\pi^P(x_3, b_2) [-(x_2 + x_3) \phi_\eta^P(x_2, b_2) + (x_2 - x_3) \phi_\eta^t(x_2, b_2)] \\
& - \phi_\pi^t(x_3, b_2) [(x_3 - x_2) \phi_\eta^P(x_2, b_2) + (x_2 + x_3) \phi_\eta^t(x_2, b_2)]] \} \\
& \cdot \alpha_s(t_f^4) h_f^4(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^4)] \} , \tag{30}
\end{aligned}$$

$$\begin{aligned}
M_{a\pi}^{P1} = & \frac{16\sqrt{6}}{3}\pi C_F m_B^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \\
& \times \{ [(x_3 - 2) r_\pi \phi_\eta(x_2, b_2) (\phi_\pi^P(x_3, b_2) + \phi_\pi^t(x_3, b_2)) - (x_2 - 2) r_\eta \phi_\pi(x_3, b_2) \\
& (\phi_\eta^P(x_2, b_2) + \phi_\eta^t(x_2, b_2))] \cdot \alpha_s(t_f^3) h_f^3(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^3)] \\
& - [x_3 r_\pi \phi_\eta(x_2, b_2) (\phi_\pi^P(x_3, b_2) + \phi_\pi^t(x_3, b_2)) \\
& - x_2 r_\eta \phi_\pi(x_3, b_2) (\phi_\eta^P(x_2, b_2) + \phi_\eta^t(x_2, b_2))] \\
& \cdot \alpha_s(t_f^4) h_f^4(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^4)] \} , \tag{31}
\end{aligned}$$

$$\begin{aligned}
M_{a\pi}^{P2} = & \frac{16\sqrt{6}}{3}\pi C_F m_B^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \\
& \times \{ [x_3 \phi_\pi(x_3, b_2) \phi_\eta(x_2, b_2) + r_\pi r_\eta (\phi_\pi^P(x_3, b_2) ((2 + x_2 + x_3) \phi_\eta^P(x_2, b_2) \\
& - (x_2 - x_3) \phi_\eta^t) + \phi_\pi^t(-(x_2 - x_3) \phi_\eta^P + (-2 + x_2 + x_3) \phi_\eta^t))] \\
& \cdot \alpha_s(t_f^3) h_f^3(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^3)] \\
& + [-x_2 \phi_\pi(x_3, b_2) \phi_\eta(x_2, b_2) - r_\pi r_\eta ((x_2 + x_3) \phi_\eta^P(x_2, b_2) \\
& - (x_2 - x_3) \phi_\eta^t(x_2, b_2) \phi_\pi^P(x_3, b_2) \\
& + ((x_3 - x_2) \phi_\eta^P(x_2, b_2) - (x_2 + x_3) \phi_\eta^t(x_2, b_2)) \phi_\pi^t(x_3, b_2))] \\
& \cdot \alpha_s(t_f^4) h_f^4(x_1, x_2, x_3, b_1, b_2) \exp[-S_{ef}(t_f^4)] \} . \tag{32}
\end{aligned}$$

In the above equations, we have assumed that $x_1 \ll x_2, x_3$. Since the light quark momentum fraction x_1 in B meson is peaked at the small region, while quark momentum fraction x_2 of $\eta^{(\prime)}$ is peaked around 0.5, this is not a bad approximation. The numerical results also show that this approximation makes very little difference in the final result. After using this approximation, all the diagrams are functions of $k_1^- = x_1 m_B / \sqrt{2}$ of B meson only, independent of the variable of k_1^+ . Therefore the integration of eq.(17) is performed safely.

If we exchange the π and $\eta^{(\prime)}$ in Fig. 1, the corresponding expressions of amplitudes for new diagrams will be similar with those as given in Eqs.(21-32), since the π and $\eta^{(\prime)}$ are all

pseudoscalar mesons and have the similar wave functions. The expressions of amplitudes for new diagrams can be obtained by the replacements

$$\phi_\pi \rightarrow \phi_\eta, \quad \phi_\pi^P \rightarrow \phi_\eta^P, \quad \phi_\pi^t \rightarrow \phi_\eta^t, \quad r_\pi \rightarrow r_\eta. \quad (33)$$

For example, we find that

$$\begin{aligned} F_{a\eta^{(\prime)}} &= -F_{a\pi}, \\ F_{a\eta^{(\prime)}}^{P1} &= -F_{a\pi}^{P1}, \\ F_{a\eta^{(\prime)}}^{P2} &= F_{a\pi}^{P2}. \end{aligned} \quad (34)$$

where the form factors $F_{a\eta^{(\prime)}}$, $F_{a\eta^{(\prime)}}^{P1}$ and $F_{a\eta^{(\prime)}}^{P2}$ describe the contributions induced by the $(V - A)(V - A)$, $(V - A)(V + A)$ and $(S - P)(S + P)$ operators, respectively.

Before we put the things together to write down the decay amplitudes for the studied decay modes, we give a brief discussion about the $\eta - \eta'$ mixing and the gluonic component of the η' meson.

The η and η' are neutral pseudoscalar ($J^P = 0^-$) mesons, and usually considered as mixtures of the $SU(3)_F$ singlet η_1 and the octet η_8 :

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \theta_p & -\sin \theta_p \\ \sin \theta_p & \cos \theta_p \end{pmatrix} \begin{pmatrix} \eta_8 \\ \eta_1 \end{pmatrix}, \quad (35)$$

with

$$\begin{aligned} \eta_8 &= \frac{1}{\sqrt{6}} (u\bar{u} + d\bar{d} - 2s\bar{s}), \\ \eta_1 &= \frac{1}{\sqrt{3}} (u\bar{u} + d\bar{d} + s\bar{s}), \end{aligned} \quad (36)$$

where θ_p is the mixing angle to be determined by various related experiments [28]. From previous studies, one obtains the mixing angle θ_p between -20° to -10° . One best fit result as given in Ref.[29] is $-17^\circ \leq \theta_p \leq -10^\circ$.

As shown in Eqs. (35,36), η and η' are generally considered as a linear combination of light quark pairs. But it should be noted that the η' meson may has a gluonic component in order to interpret the anomalously large branching ratios of $B \rightarrow K\eta'$ and $J/\Psi \rightarrow \eta'\gamma$ [29, 30]. In Refs.[29, 30, 31], the physical states η and η' were defined as

$$\begin{aligned} |\eta\rangle &= X_\eta \left| \frac{u\bar{u} + d\bar{d}}{\sqrt{2}} \right\rangle + Y_\eta |s\bar{s}\rangle, \\ |\eta'\rangle &= X_{\eta'} \left| \frac{u\bar{u} + d\bar{d}}{\sqrt{2}} \right\rangle + Y_{\eta'} |s\bar{s}\rangle + Z_{\eta'} |gluonium\rangle, \end{aligned} \quad (37)$$

where $X_{\eta^{(\prime)}}$, $Y_{\eta^{(\prime)}}$ and $Z_{\eta'}$ parameters describe the ratios of $u\bar{u} + d\bar{d}$, $s\bar{s}$ and gluonium ($SU(3)_F$ singlet) componetnt of $\eta^{(\prime)}$, respectively. In Ref.[29], the author shows that the gluonic admixture in η' can be as large as 26%, i.e.

$$Z_{\eta'} / (X_{\eta^{(\prime)}} + Y_{\eta^{(\prime)}} + Z_{\eta'}) \leq 0.26. \quad (38)$$

According to paper [30], a large SU(3) singlet contribution can help us to explain the large branching ratio for $B \rightarrow K\eta'$ decay, but also result in a large branching ratio for $B \rightarrow K^0\eta$ decay, $Br(B \rightarrow K^0\eta) \sim 7.0(13) \times 10^{-6}$ for $\theta_P = -20^\circ(-10^\circ)$ as given in Table II of Ref.[30], which is clearly too large than currently available upper limits [19]:

$$Br(B \rightarrow K^0\eta) < 1.9 \times 10^{-6}. \quad (39)$$

Although a lot of studies have been done along this direction, but we currently still do not understand the anomalous $gg - \eta'$ coupling clearly, and do not know how to calculate reliably the contributions induced by the gluonic component of η' meson. In this paper, we firstly assume that η' does not have the gluonic component, and set the quark content of η and η' as described by Eqs. (35,36). We will also discuss the effects of a non-zero gluonic admixture of η' in next section.

Combining the contributions from different diagrams, the total decay amplitude for $B^+ \rightarrow \pi^+\eta$ decay can be written as

$$\begin{aligned} \sqrt{3}\mathcal{M}(\pi^+\eta) = & F_{e\pi} \left\{ \left[\xi_u \left(C_1 + \frac{1}{3}C_2 \right) \right. \right. \\ & - \xi_t \left(\frac{7}{3}C_3 + \frac{5}{3}C_4 - 2C_5 - \frac{2}{3}C_6 - \frac{1}{2}C_7 - \frac{1}{6}C_8 + \frac{1}{3}C_9 - \frac{1}{3}C_{10} \right) \left. \right] f_\eta^d F_1(\theta_p) \\ & - \xi_t \left(C_3 + \frac{1}{3}C_4 - C_5 - \frac{1}{3}C_6 + \frac{1}{2}C_7 + \frac{1}{6}C_8 - \frac{1}{2}C_9 - \frac{1}{6}C_{10} \right) f_\eta^s F_2(\theta_p) \left. \right\} \\ & - F_{e\pi}^{P_2} \xi_t \left(\frac{1}{3}C_5 + C_6 - \frac{1}{6}C_7 - \frac{1}{2}C_8 \right) f_\eta^d F_1(\theta_p) \\ & + M_{e\pi} \left\{ \left[\xi_u C_2 - \xi_t \cdot \left(C_3 + 2C_4 - \frac{1}{2}C_9 + \frac{1}{2}C_{10} \right) \right] F_1(\theta_p) \right. \\ & - \xi_t \left(C_4 - \frac{1}{2}C_{10} \right) F_2(\theta_p) \left. \right\} \\ & - M_{e\pi}^{P_2} \xi_t \left[\left(2C_6 + \frac{1}{2}C_8 \right) F_1(\theta_p) + \left(C_6 - \frac{1}{2}C_8 \right) F_2(\theta_p) \right] \\ & + (M_{a\pi} + M_e + M_a) [\xi_u C_1 - \xi_t (C_3 + C_9)] F_1(\theta_p) \\ & - (M_{a\pi}^{P_1} + M_a^{P_1}) \xi_t (C_5 + C_7) \cdot F_1(\theta_p) \\ & + (F_{a\pi}^{P_2} + F_e^{P_2} + F_a^{P_2}) (-\xi_t) \left(C_6 + C_8 + \frac{1}{3}C_5 + \frac{1}{3}C_7 \right) F_1(\theta_p) \\ & + F_e \cdot \left\{ \left[\xi_u \left(\frac{1}{3}C_1 + C_2 \right) - \xi_t \left(\frac{1}{3}C_3 + C_4 + \frac{1}{3}C_9 + C_{10} \right) \right] F_1(\theta_p) \right\}, \quad (40) \end{aligned}$$

where $\xi_u = V_{ub}^* V_{ud}$, $\xi_t = V_{tb}^* V_{td}$, and

$$\begin{aligned} F_1(\theta_p) &= -\sin \theta_p + \cos \theta_p / \sqrt{2}, \\ F_2(\theta_p) &= -\sin \theta_p - \sqrt{2} \cos \theta_p, \end{aligned} \quad (41)$$

are the mixing factors. The Wilson coefficients C_i should be calculated at the appropriate scale t using equations as given in the Appendices of Ref. [9].

Similarly, the decay amplitude for $B^0 \rightarrow \pi^0 \eta$ can be written as

$$\begin{aligned}
\sqrt{6}\mathcal{M}(\pi^0\eta) = & F_e \left[\xi_u \left(C_1 + \frac{1}{3}C_2 \right) - \xi_t \left(-\frac{1}{3}C_3 - C_4 - \frac{3}{2}C_7 - \frac{1}{2}C_8 + \frac{5}{3}C_9 + C_{10} \right) \right] F_1(\theta_p) \\
& - F_{e\pi} \left[\xi_u \left(C_1 + \frac{1}{3}C_2 \right) \cdot f_\eta^d F_1(\theta_p) \right. \\
& - \xi_t \left(\frac{7}{3}C_3 + \frac{5}{3}C_4 - 2C_5 - \frac{2}{3}C_6 - \frac{1}{2}C_7 - \frac{1}{6}C_8 + \frac{1}{3}C_9 - \frac{1}{3}C_{10} \right) \cdot f_\eta^d F_1(\theta_p) \\
& \left. - \xi_t \left(C_3 + \frac{1}{3}C_4 - C_5 - \frac{1}{3}C_6 + \frac{1}{2}C_7 + \frac{1}{6}C_8 - \frac{1}{2}C_9 - \frac{1}{6}C_{10} \right) \cdot f_\eta^s F_2(\theta_p) \right] \\
& - M_{e\pi} \left\{ \left[\xi_u C_2 - \xi_t \left(C_3 + 2C_4 - \frac{1}{2}C_9 + \frac{1}{2}C_{10} \right) \right] \cdot F_1(\theta_p) \right. \\
& \left. - \xi_t \left(C_4 - \frac{1}{2}C_{10} \right) F_2(\theta_p) \right\} \\
& - M_{e\pi}^{P_2} \left[-\xi_t \left(2C_6 + \frac{1}{2}C_8 \right) F_1(\theta_p) - \xi_t \left(C_6 - \frac{1}{2}C_8 \right) F_2(\theta_p) \right] \\
& - (F_{a\pi}^{P_2} + F_a^{P_2} + F_e^{P_2}) \left[-\xi_t \left(C_6 + \frac{1}{3}C_5 - \frac{1}{2}C_8 - \frac{1}{6}C_7 \right) \right] \cdot F_1(\theta_p) \\
& + (M_{a\pi} + M_a + M_e) \left[\xi_u C_2 - \xi_t \left(-C_3 + \frac{1}{2}C_9 + \frac{3}{2}C_{10} \right) \right] F_1(\theta_p) \\
& + (M_a^{P_1} + M_{a\pi}^{P_1}) \xi_t \left(C_5 - \frac{1}{2}C_7 \right) F_1(\theta_p) \\
& - \frac{3}{2} (M_{a\pi}^{P_2} + M_a^{P_2} + M_e^{P_2}) \xi_t C_8 F_1(\theta_p) \\
& + F_{e\pi}^{P_2} \xi_t \left(\frac{1}{3}C_5 + C_6 - \frac{1}{6}C_7 - \frac{1}{2}C_8 \right) \cdot F_1(\theta_p). \tag{42}
\end{aligned}$$

The decay amplitudes for $B \rightarrow \pi^+ \eta'$ and $B \rightarrow \pi^0 \eta'$ can be obtained easily from Eqs.(40) and (42) by the following replacements

$$\begin{aligned}
f_\eta^d, f_\eta^s & \longrightarrow f_{\eta'}^d, f_{\eta'}^s, \\
F_1(\theta_p) & \longrightarrow F_1'(\theta_p) = \cos \theta_p + \frac{\sin \theta_p}{\sqrt{2}}, \\
F_2(\theta_p) & \longrightarrow F_2'(\theta_p) = \cos \theta_p - \sqrt{2} \sin \theta_p. \tag{43}
\end{aligned}$$

Note that the possible gluonic component of η' meson has been neglected here.

IV. NUMERICAL RESULTS AND DISCUSSIONS

A. Input parameters and wave functions

We use the following input parameters in the numerical calculations

$$\begin{aligned}\Lambda_{\overline{\text{MS}}}^{(f=4)} &= 250\text{MeV}, \quad f_\pi = 130\text{MeV}, \quad f_B = 190\text{MeV}, \\ m_0^{\eta_{d\bar{d}}} &= 1.4\text{GeV}, \quad m_0^{\eta_{s\bar{s}}} = 2.4\text{GeV}, \quad f_K = 160\text{MeV}, \\ M_B &= 5.2792\text{GeV}, \quad M_W = 80.41\text{GeV}.\end{aligned}\tag{44}$$

For the CKM matrix elements, here we adopt the wolfenstein parametrization for the CKM matrix up to $\mathcal{O}(\lambda^3)$,

$$V_{CKM} = \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix},\tag{45}$$

with the parameters $\lambda = 0.22$, $A = 0.853$, $\rho = 0.20$ and $\eta = 0.33$.

For the B meson wave function, we adopt the model

$$\phi_B(x, b) = N_B x^2 (1-x)^2 \exp \left[-\frac{M_B^2 x^2}{2\omega_b^2} - \frac{1}{2}(\omega_b b)^2 \right],\tag{46}$$

where ω_b is a free parameter and we take $\omega_b = 0.4 \pm 0.05$ GeV in numerical calculations, and $N_B = 91.745$ is the normalization factor for $\omega_b = 0.4$. This is the same wave functions as in Refs.[9, 10, 25, 26], which is a best fit for most of the measured hadronic B decays.

For the light meson wave function, we neglect the b dependant part, which is not important in numerical analysis. We choose the wave function of π meson [32]:

$$\phi_\pi^A(x) = \frac{3}{\sqrt{6}} f_\pi x(1-x) \left[1 + 0.44 C_2^{3/2}(2x-1) + 0.25 C_4^{3/2}(2x-1) \right],\tag{47}$$

$$\phi_\pi^P(x) = \frac{f_\pi}{2\sqrt{6}} \left[1 + 0.43 C_2^{1/2}(2x-1) + 0.09 C_4^{1/2}(2x-1) \right],\tag{48}$$

$$\phi_\pi^t(x) = \frac{f_\pi}{2\sqrt{6}} (1-2x) \left[1 + 0.55(10x^2 - 10x + 1) \right].\tag{49}$$

The Gegenbauer polynomials are defined by

$$\begin{aligned}C_2^{1/2}(t) &= \frac{1}{2}(3t^2 - 1), \quad C_4^{1/2}(t) = \frac{1}{8}(35t^4 - 30t^2 + 3), \\ C_2^{3/2}(t) &= \frac{3}{2}(5t^2 - 1), \quad C_4^{3/2}(t) = \frac{15}{8}(21t^4 - 14t^2 + 1).\end{aligned}\tag{50}$$

For η meson's wave function, $\phi_{\eta_{d\bar{d}}}^A$, $\phi_{\eta_{d\bar{d}}}^P$ and $\phi_{\eta_{d\bar{d}}}^T$ represent the axial vector, pseudoscalar and tensor components of the wave function respectively, for which we utilize the result

from the light-cone sum rule [33] including twist-3 contribution:

$$\begin{aligned}
\phi_{\eta_{d\bar{d}}}^A(x) &= \frac{3}{\sqrt{2N_c}} f_x x(1-x) \left\{ 1 + a_2^{\eta_{d\bar{d}}} \frac{3}{2} [5(1-2x)^2 - 1] \right. \\
&\quad \left. + a_4^{\eta_{d\bar{d}}} \frac{15}{8} [21(1-2x)^4 - 14(1-2x)^2 + 1] \right\}, \\
\phi_{\eta_{d\bar{d}}}^P(x) &= \frac{1}{2\sqrt{2N_c}} f_x \left\{ 1 + \frac{1}{2} \left(30\eta_3 - \frac{5}{2}\rho_{\eta_{d\bar{d}}}^2 \right) [3(1-2x)^2 - 1] \right. \\
&\quad \left. + \frac{1}{8} \left(-3\eta_3\omega_3 - \frac{27}{20}\rho_{\eta_{d\bar{d}}}^2 - \frac{81}{10}\rho_{\eta_{d\bar{d}}}^2 a_2^{\eta_{d\bar{d}}} \right) [35(1-2x)^4 - 30(1-2x)^2 + 3] \right\}, \\
\phi_{\eta_{d\bar{d}}}^T(x) &= \frac{3}{\sqrt{2N_c}} f_x (1-2x) \\
&\quad \cdot \left[\frac{1}{6} + (5\eta_3 - \frac{1}{2}\eta_3\omega_3 - \frac{7}{20}\rho_{\eta_{d\bar{d}}}^2 - \frac{3}{5}\rho_{\eta_{d\bar{d}}}^2 a_2^{\eta_{d\bar{d}}})(10x^2 - 10x + 1) \right],
\end{aligned} \tag{51}$$

with

$$\begin{aligned}
a_2^{\eta_{d\bar{d}}} &= 0.44, \quad a_4^{\eta_{d\bar{d}}} = 0.25, \quad a_1^K = 0.20, \quad a_2^K = 0.25, \\
\rho_{\eta_{d\bar{d}}} &= m_\pi/m_0^{\eta_{d\bar{d}}}, \quad \eta_3 = 0.015, \quad \omega_3 = -3.0.
\end{aligned} \tag{52}$$

We assume that the wave function of $u\bar{u}$ is same as the wave function of $d\bar{d}$. For the wave function of the $s\bar{s}$ components, we also use the same form as $d\bar{d}$ but with $m_0^{s\bar{s}}$ and f_y instead of $m_0^{d\bar{d}}$ and f_x , respectively. For f_x and f_y , we use the values as given in Ref.[34] where isospin symmetry is assumed for f_x and $SU(3)$ breaking effect is included for f_y :

$$f_x = f_\pi, \quad f_y = \sqrt{2f_K^2 - f_\pi^2}. \tag{53}$$

These values are translated to the values in the two mixing angle method, which is often used in vacuum saturation approach as:

$$\begin{aligned}
f_8 &= 169\text{MeV}, \quad f_1 = 151\text{MeV}, \\
\theta_8 &= -25.9^\circ(-18.9^\circ), \quad \theta_1 = -7.1^\circ(-0.1^\circ),
\end{aligned} \tag{54}$$

where the pseudoscalar mixing angle θ_p is taken as -17° (-10°) [29]. The parameters m_0^i ($i = \eta_{d\bar{d}(u\bar{u})}, \eta_{s\bar{s}}$) are defined as:

$$m_0^{\eta_{d\bar{d}(u\bar{u})}} \equiv m_0^\pi \equiv \frac{m_\pi^2}{(m_u + m_d)}, \quad m_0^{\eta_{s\bar{s}}} \equiv \frac{2M_K^2 - m_\pi^2}{(2m_s)}. \tag{55}$$

We include full expression of twist-3 wave functions for light mesons. The twist-3 wave functions are also adopted from QCD sum rule calculations [35]. We will see later that this set of parameters will give good results for $B \rightarrow \pi\eta^{(\prime)}$ decays. Using the above chosen wave functions and the central values of relevant input parameters, we find the numerical values of the corresponding form factors at zero momentum transfer:

$$\begin{aligned}
F_0^{B \rightarrow \pi}(q^2 = 0) &= 0.30, \\
F_0^{B \rightarrow \eta}(q^2 = 0) &= 0.15, \\
F_0^{B \rightarrow \eta'}(q^2 = 0) &= 0.14.
\end{aligned} \tag{56}$$

These values agree well with those as given in Refs. [33, 34, 36].

B. Branching ratios

For $B \rightarrow \pi\eta^{(\prime)}$ decays, the decay amplitudes in Eqs. (40) and (42) can be rewritten as

$$\mathcal{M} = V_{ub}^* V_{ud} T - V_{tb}^* V_{td} P = V_{ub}^* V_{ud} T [1 + z e^{i(\alpha+\delta)}], \quad (57)$$

where

$$z = \left| \frac{V_{tb}^* V_{td}}{V_{ub}^* V_{ud}} \right| \left| \frac{P}{T} \right| \quad (58)$$

is the ratio of penguin to tree contributions, $\alpha = \arg \left[-\frac{V_{td} V_{tb}^*}{V_{ud} V_{ub}^*} \right]$ is the weak phase (one of the three CKM angles), and δ is the relative strong phase between tree (T) and penguin (P) diagrams. The ratio z and the strong phase δ can be calculated in the pQCD approach. One can leave the CKM angle α as a free parameter and explore the CP asymmetry parameter dependence on it.

For $B \rightarrow \pi^+ \eta$ decay, for example, one can find “T” and “P” terms by comparing the decay amplitude as defined in Eq. (40) with that in Eq. (57),

$$\begin{aligned} T(\pi^+ \eta) = & \frac{F_1(\theta_p)}{\sqrt{3}} \cdot \left\{ F_{e\pi} \left(C_1 + \frac{1}{3} C_2 \right) f_\eta^d + M_{e\pi} C_2 \right. \\ & \left. + F_e \left(\frac{1}{3} C_1 + C_2 \right) + (M_a + M_e + M_{a\pi}) C_1 \right\}, \end{aligned} \quad (59)$$

$$\begin{aligned} \sqrt{3} P(\pi^+ \eta) = & F_{e\pi} \left[\left(\frac{7}{3} C_3 + \frac{5}{3} C_4 - 2C_5 - \frac{2}{3} C_6 - \frac{1}{2} C_7 - \frac{1}{6} C_8 + \frac{1}{3} C_9 - \frac{1}{3} C_{10} \right) f_\eta^d F_1(\theta_p) \right. \\ & \left. + \left(C_3 + \frac{1}{3} C_4 - C_5 - \frac{1}{3} C_6 + \frac{1}{2} C_7 + \frac{1}{6} C_8 - \frac{1}{2} C_9 - \frac{1}{6} C_{10} \right) f_\eta^s F_2(\theta_p) \right] \\ & + F_{e\pi}^{P_2} \left(\frac{1}{3} C_5 + C_6 - \frac{1}{6} C_7 - \frac{1}{2} C_8 \right) f_\eta^d F_1(\theta_p) \\ & - M_{e\pi} \left[- \left(C_3 + 2C_4 - \frac{1}{2} C_9 + \frac{1}{2} C_{10} \right) F_1(\theta_p) - \left(C_4 - \frac{1}{2} C_{10} \right) F_2(\theta_p) \right] \\ & + M_{e\pi}^{P_2} \left[\left(2C_6 + \frac{1}{2} C_8 \right) F_1(\theta_p) + \left(C_6 - \frac{1}{2} C_8 \right) F_2(\theta_p) \right] \\ & + (M_{a\pi} + M_e + M_a)(C_3 + C_9) F_1(\theta_p) + (M_{a\pi}^{P_1} + M_a^{P_1})(C_5 + C_7) \cdot F_1(\theta_p) \\ & + (F_{a\pi}^{P_2} + F_e^{P_2} + F_a^{P_2}) F_1(\theta_p) \left(C_6 + C_8 + \frac{1}{3} C_5 + \frac{1}{3} C_7 \right) \\ & + F_e \left(\frac{1}{3} C_3 + C_4 + \frac{1}{3} C_9 + C_{10} \right) F_1(\theta_p), \end{aligned} \quad (60)$$

Similarly, one can obtain the expressions of the corresponding tree and penguin terms for the remaining three decays.

Using the “T” and “P” terms as given in Eqs.(59) and (60), it is easy to calculate the ratio z and the strong phase δ for the decay in study. For $B^+ \rightarrow \rho^+ \eta$ and $\rho^+ \eta'$ decays, we find numerically that

$$z(\pi^+ \eta) = 0.33, \quad \delta(\pi^+ \eta) = -136^\circ, \quad (61)$$

$$z(\pi^+ \eta') = 0.25, \quad \delta(\pi^+ \eta') = -130^\circ. \quad (62)$$

The main errors of the ratio z and the strong phase δ are induced by the uncertainty of $\omega_b = 0.4 \pm 0.05$ GeV and $m_0^\pi = 1.4 \pm 0.1$ GeV and small in magnitude. The reason is that the errors induced by the uncertainties of input parameters are largely cancelled in the ratio. We therefore use the central values of z and δ in the following numerical calculations, unless explicitly stated otherwise.

From Eq. (57), it is easy to write the decay amplitude for the corresponding charge conjugated decay mode

$$\overline{\mathcal{M}} = V_{ub}V_{ud}^*T - V_{tb}V_{td}^*P = V_{ub}V_{ud}^*T [1 + ze^{i(-\alpha+\delta)}]. \quad (63)$$

Therefore the CP-averaged branching ratio for $B^0 \rightarrow \pi \eta^{(\prime)}$ is

$$Br = (|\mathcal{M}|^2 + |\overline{\mathcal{M}}|^2)/2 = |V_{ub}V_{ud}^*T|^2 [1 + 2z \cos \alpha \cos \delta + z^2], \quad (64)$$

where the ratio z and the strong phase δ have been defined in Eqs.(57) and (58). It is easy to see that the CP-averaged branching ratio is a function of $\cos \alpha$. This gives a potential method to determine the CKM angle α by measuring only the CP-averaged branching ratios with PQCD calculations.

Using the wave functions and the input parameters as specified in previous sections, it is straightforward to calculate the branching ratios for the four considered decays. The theoretical predictions in the PQCD approach for the branching ratios of the decays under consideration are the following

$$Br(B^+ \rightarrow \pi^+ \eta) = [4.1_{-0.9}^{+1.3}(\omega_b)_{-0.3}^{+0.4}(m_0^\pi)_{-0.5}^{+0.6}(\alpha)] \times 10^{-6}, \quad (65)$$

$$Br(B^+ \rightarrow \pi^+ \eta') = [2.4_{-0.5}^{+0.8}(\omega_b) \pm 0.2(m_0^\pi) \pm 0.3(\alpha)] \times 10^{-6}, \quad (66)$$

$$Br(B^0 \rightarrow \pi^0 \eta) = [0.23_{-0.03}^{+0.04}(\omega_b)_{-0.03}^{+0.04}(m_0^\pi) \pm 0.05(\alpha)] \times 10^{-6}, \quad (67)$$

$$Br(B^0 \rightarrow \pi^0 \eta') = [0.19 \pm 0.02(\omega_b) \pm 0.03(m_0^\pi)_{-0.05}^{+0.04}(\alpha)] \times 10^{-6}, \quad (68)$$

for $\theta_p = -17^\circ$, and

$$Br(B^+ \rightarrow \pi^+ \eta) = [3.3_{-0.8}^{+1.0}(\omega_b) \pm 0.3(m_0^\pi) \pm 0.4(\alpha)] \times 10^{-6}, \quad (69)$$

$$Br(B^+ \rightarrow \pi^+ \eta') = [3.2_{-0.7}^{+1.1}(\omega_b) \pm 0.3(m_0^\pi) \pm 0.4(\alpha)] \times 10^{-6}, \quad (70)$$

$$Br(B^0 \rightarrow \pi^0 \eta) = [0.17 \pm 0.02(\omega_b) \pm 0.02(m_0^\pi)_{-0.04}^{+0.03}(\alpha)] \times 10^{-6}, \quad (71)$$

$$Br(B^0 \rightarrow \pi^0 \eta') = [0.28_{-0.02}^{+0.04}(\omega_b) \pm 0.04(m_0^\pi) \pm 0.05(\alpha)] \times 10^{-6}, \quad (72)$$

for $\theta_p = -10^\circ$. The main errors are induced by the uncertainties of $\omega_b = 0.4 \pm 0.04$ GeV, $m_0^\pi = 1.4 \pm 0.1$ GeV and $\alpha = 100^\circ \pm 20^\circ$, respectively.

The PQCD predictions for the branching ratios of considered decays agree well with the measured values or the upper limits as shown in Eqs.(1-5). Furthermore, the pQCD




FIG. 2: The α dependence of the branching ratios (in unit of 10^{-6}) of $B^+ \rightarrow \pi^+ \eta^{(\prime)}$ decays for $m_0^\pi = 1.4$ GeV, $\theta_p = -17^\circ$, $\omega_b = 0.36$ GeV (dotted curve), 0.40 GeV (solid curve) and 0.44 GeV (dashed curve). The gray band shows the experimental data.

predictions also agree well with the theoretical predictions in the QCDF approach, for example, as given in Ref. [8]:

$$\begin{aligned}
 Br(B^+ \rightarrow \pi^+ \eta) &= (4.7_{-2.3}^{+2.7}) \times 10^{-6}, \\
 Br(B^+ \rightarrow \pi^+ \eta') &= (3.1_{-1.7}^{+1.9}) \times 10^{-6}, \\
 Br(B^0 \rightarrow \pi^0 \eta) &= (0.28_{-0.28}^{+0.48}) \times 10^{-6}, \\
 Br(B^0 \rightarrow \pi^0 \eta') &= (0.17_{-0.17}^{+0.33}) \times 10^{-6},
 \end{aligned} \tag{73}$$

where the individual errors as given in Ref. [8] have been added in quadrature.

It is worth stressing that the theoretical predictions in the PQCD approach have relatively large theoretical errors induced by the still large uncertainties of many input parameters, such as ω_b , m_0^π and θ_p . In our analysis, we consider the constraints on these parameters from analysis of other well measured decay channels. For example, the constraint $1.1\text{GeV} \leq m_0^\pi \leq 1.9\text{GeV}$ was obtained from the phenomenological studies for $B \rightarrow \pi\pi$ decays [9], while the constraint of $\alpha \approx 100^\circ \pm 20^\circ$ were obtained by direct measurements or from the global fit [19, 37]. From numerical calculations, we get to know that the main errors come from the uncertainty of ω_b , m_0^π , α and θ_p .

In Figs. 2 and 3, we present, respectively, the pQCD predictions of the branching ratios of $B \rightarrow \pi^+ \eta$ and $\pi^+ \eta'$ decays for $\theta_p = 10^\circ$, $\omega_b = 0.4 \pm 0.04$ GeV, $m_0^\pi = 1.4 \pm 0.1$ GeV and $\alpha = [0^\circ, 180^\circ]$. Fig. 4 shows the α -dependence of the pQCD predictions of the branching ratios of $B \rightarrow \pi^0 \eta^{(\prime)}$ decays for $\theta_p = 10^\circ$, $\omega_b = 0.4$ GeV, $m_0^\pi = 1.4$ GeV and $\alpha = [0^\circ, 180^\circ]$.

From the numerical results and the figures we observe that the pQCD predictions are sensitive to the variations of ω_b and m_0^π . The parameter m_0^π originates from the chiral perturbation theory and have a value near 1 GeV. The m_0^π parameter characterizes the relative size of twist 3 contribution to twist 2 contribution. Because of the chiral enhancement of m_0^π , the twist 3 contribution become comparable in size with the twist 2 contribution. The branching ratios of $Br(B \rightarrow \pi \eta^{(\prime)})$ are also sensitive to the parameter




FIG. 3: The α dependence of the branching ratios (in unit of 10^{-6}) of $B^+ \rightarrow \pi^+ \eta^{(\prime)}$ decays for $\omega_b = 0.4$ GeV, $\theta_p = -17^\circ$, $m_0^\pi = 1.3$ GeV (dotted curve), 1.4 GeV (solid curve) and 1.5 GeV (dashed curve). The gray band shows the data.




FIG. 4: The α dependence of the branching ratios (in unit of 10^{-6}) of $\pi^0 \eta$ (solid curve) and $\pi^0 \eta'$ (dotted curve) decays for $m_0^\pi = 1.4$ GeV, $\theta_p = -17^\circ$, $\omega_b = 0.40$ GeV.

m_0^π , but not as strong as the ω_b dependence.

C. CP-violating asymmetries

Now we turn to the evaluations of the CP-violating asymmetries of $B \rightarrow \pi \eta^{(\prime)}$ decays in pQCD approach. For $B^+ \rightarrow \pi^+ \eta$ and $B^+ \rightarrow \pi^+ \eta'$ decays, the direct CP-violating asymmetries $\mathcal{A}_{CP}$ can be defined as:

$$\mathcal{A}_{CP}^{dir} = \frac{|\overline{\mathcal{M}}|^2 - |\mathcal{M}|^2}{|\overline{\mathcal{M}}|^2 + |\mathcal{M}|^2} = \frac{2z \sin \alpha \sin \delta}{1 + 2z \cos \alpha \cos \delta + z^2}, \quad (74)$$

where the ratio z and the strong phase δ have been defined in previous subsection and are calculable in pQCD approach.

It is easy to calculate the CP-violating asymmetries with z and δ . In Fig. 5, we show the α -dependence of the direct CP-violating asymmetries $\mathcal{A}_{CP}^{dir}$ for $B^\pm \rightarrow \pi^\pm \eta$ (the solid curve) and $B^\pm \rightarrow \pi^\pm \eta'$ (the dotted curve) decay, respectively. From Fig. 5, one can see that the CP-violating asymmetries $\mathcal{A}_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta^{(\prime)})$ are large in magnitude, about 35% for $\alpha \sim 100^\circ$. So large CP-violating asymmetry plus large ($\sim 10^{-6}$) branching ratios are measurable in current B factory experiments.

The pQCD predictions for $\mathcal{A}_{CP}^{dir}$ and the major theoretical errors for $B^\pm \rightarrow \pi^\pm \eta^{(\prime)}$ decays are

$$\mathcal{A}_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta) = (-37_{-6}^{+8}(\alpha) \pm 4(\omega_b)_{-1}^{+0}(m_0^\pi)) \times 10^{-2}, \quad (75)$$

$$\mathcal{A}_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta') = (-33_{-4}^{+6}(\alpha)_{-6}^{+4}(\omega_b)_{-2}^{+0}(m_0^\pi)) \times 10^{-2}, \quad (76)$$

where the dominant errors come from the variations of $\omega_b = 0.4 \pm 0.04$ GeV, $m_0^\pi = 1.4 \pm 0.1$ GeV and $\alpha = 100^\circ \pm 20^\circ$.

By comparing the above numerical results with those measured values as given in Eqs.(3) and (4)¹, we find that

1. The pQCD predictions for the direct CP-violating asymmetry for both $B \rightarrow \pi\eta$ and $\pi\eta'$ decays are large in magnitude and have a moderate theoretical error because of the cancelation in the ratios.
2. For $B^\pm \rightarrow \pi^\pm \eta$ decay, the pQCD prediction for $\mathcal{A}_{CP}^{dir}$ has the same sign with the measured value and also consistent with it within 2σ errors. For $B^\pm \rightarrow \pi^\pm \eta'$ decay, however, the pQCD prediction for $\mathcal{A}_{CP}^{dir}$ has the opposite sign with the measured value.
3. For $B^\pm \rightarrow \pi^\pm \eta'$ decay, although there exist a clear difference between the pQCD prediction of $\mathcal{A}_{CP}^{dir}$ and the data, but it is too early to draw any reliable information from such difference because of the still large theoretical and experimental errors. More theoretical studies (for example, calculation of next-to-leading order contributions [39]) and more accurate measurements are needed to clarify this discrepancy.

In Ref. [8], by using the “default values” of input parameters, the authors presented their predictions for $\mathcal{A}_{CP}^{dir}(B \rightarrow \pi^\pm \eta^{(\prime)})$ in the QCDF approach

$$\mathcal{A}_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta) = (-14.9_{-5.4}^{+4.9} {}_{-7.4}^{+8.3} {}_{-0.8}^{+1.3} {}_{-17.3}^{+17.4}) \times 10^{-2}, \quad (77)$$

$$\mathcal{A}_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta') = (-8.6_{-3.1}^{+2.8} {}_{-9.0}^{+10.5} {}_{-0.7}^{+0.7} {}_{-20.4}^{+20.4}) \times 10^{-2}, \quad (78)$$

where the first error comes from the variation of the CKM parameters, the second error refers to the variation of $\mu \sim m_b$, quark masses, decay constants, form factors, and the mixing angle θ_p . The third error corresponds to the uncertainty due to the Gegenbauer moments in the expansion of the LCDAs. The last but largest error is induced by the

¹ We used the same convention as the BaBar and Belle Collaboration [17, 18] to define the CP-violating asymmetry for $B \rightarrow \pi\eta^{(\prime)}$ decays.




FIG. 5: The direct CP asymmetries (in percentage) of $B^+ \rightarrow \pi^+ \eta$ (solid curve) and $B^+ \rightarrow \pi^+ \eta'$ (dotted curve) as a function of CKM angle α .

uncertainty of the unknown annihilation contributions. In fact, these numbers are similar to those generalized factorization approach [38], since the mechanism of strong phase is the same for these two approaches.

By using the “set S_4 ” input parameters [8], the central values of the QCDF prediction can become positive simultaneously

$$\mathcal{A}_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta) = 5.6 \times 10^{-2}, \quad (79)$$

$$\mathcal{A}_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta') = 11.1 \times 10^{-2}. \quad (80)$$

From the numerical results as given in Eqs.(77-80), one can see that the dominant theoretical error from the annihilation contributions in QCD factorization approach are too large to make any meaningful comparisons between the theoretical predictions and the data for $\mathcal{A}_{CP}^{dir}(B \rightarrow \pi \eta^{(\prime)})$. The reason is that the annihilation contributions play a key role for producing the strong phase of the two-body charmless B meson decays, but unfortunately they are incalculable in QCD factorization approach.

We now study the CP-violating asymmetries for $B^0 \rightarrow \pi^0 \eta^{(\prime)}$ decays. For these neutral decay modes, the effects of $B^0 - \bar{B}^0$ mixing should be considered. For B^0 meson decays, we know that $\Delta\Gamma/\Delta m_d \ll 1$ and $\Delta\Gamma/\Gamma \ll 1$. The CP-violating asymmetry of $B^0(\bar{B}^0) \rightarrow \pi^0 \eta^{(\prime)}$ decay is time dependent and can be defined as

$$\begin{aligned} A_{CP} &\equiv \frac{\Gamma(\bar{B}_d^0(\Delta t) \rightarrow f_{CP}) - \Gamma(B_d^0(\Delta t) \rightarrow f_{CP})}{\Gamma(\bar{B}_d^0(\Delta t) \rightarrow f_{CP}) + \Gamma(B_d^0(\Delta t) \rightarrow f_{CP})} \\ &= A_{CP}^{dir} \cos(\Delta m \Delta t) + A_{CP}^{mix} \sin(\Delta m \Delta t), \end{aligned} \quad (81)$$

where Δm is the mass difference between the two B_d^0 mass eigenstates, $\Delta t = t_{CP} - t_{tag}$ is the time difference between the tagged B^0 ($\bar{B}^0$) and the accompanying $\bar{B}^0$ (B^0) with opposite b flavor decaying to the final CP-eigenstate f_{CP} at the time t_{CP} . The direct and

mixing induced CP-violating asymmetries A_{CP}^{dir} and A_{CP}^{mix} can be written as

$$\mathcal{A}_{CP}^{dir} = \frac{|\lambda_{CP}|^2 - 1}{1 + |\lambda_{CP}|^2}, \quad A_{CP}^{mix} = \frac{2Im(\lambda_{CP})}{1 + |\lambda_{CP}|^2}, \quad (82)$$

where the CP-violating parameter λ_{CP} is

$$\lambda_{CP} = \frac{V_{tb}^* V_{td} \langle \pi^0 \eta^{(\prime)} | H_{eff} | \overline{B}^0 \rangle}{V_{tb} V_{td}^* \langle \pi^0 \eta^{(\prime)} | H_{eff} | B^0 \rangle} = e^{2i\alpha} \frac{1 + ze^{i(\delta-\alpha)}}{1 + ze^{i(\delta+\alpha)}}. \quad (83)$$

Here the ratio z and the strong phase δ have been defined previously. In PQCD approach, since both z and δ are calculable, it is easy to find the numerical values of A_{CP}^{dir} and A_{CP}^{mix} for the considered decay processes.

In Figs. 6 and 7, we show the α -dependence of the direct and the mixing-induced CP-violating asymmetry A_{CP}^{dir} and A_{CP}^{mix} for $B^0 \rightarrow \pi^0 \eta$ (solid curve) and $B^0 \rightarrow \pi^0 \eta'$ (dotted curve) decays, respectively.

The pQCD predictions for the direct and mixing induced CP-violating asymmetries of $B^0 \rightarrow \pi^0 \eta^{(\prime)}$ decays are

$$\mathcal{A}_{CP}^{dir}(B^0 \rightarrow \pi^0 \eta) = (-42_{-12}^{+9}(\alpha)_{-2}^{+3}(\omega_b)_{-3}^{+1}(m_0^\pi)) \times 10^{-2}, \quad (84)$$

$$\mathcal{A}_{CP}^{dir}(B^0 \rightarrow \pi^0 \eta') = (-36_{-9}^{+10}(\alpha)_{-1}^{+2}(\omega_b)_{-3}^{+2}(m_0^\pi)) \times 10^{-2}, \quad (85)$$

$$\mathcal{A}_{CP}^{mix}(B^0 \rightarrow \pi^0 \eta) = (67_{-9}^{+0}(\alpha)_{-6}^{+5}(\omega_b)_{-2}^{+1}(m_0^\pi)) \times 10^{-2}, \quad (86)$$

$$\mathcal{A}_{CP}^{mix}(B^0 \rightarrow \pi^0 \eta') = (67_{-9}^{+0}(\alpha)_{-6}^{+4}(\omega_b)_{-3}^{+1}(m_0^\pi)) \times 10^{-2}, \quad (87)$$

where the dominant errors come from the variations of $\omega_b = 0.4 \pm 0.05$ GeV, $m_0^\pi = 1.4 \pm 0.1$ GeV and $\alpha = 100^\circ \pm 20^\circ$.

As a comparison, we present the QCDF predictions for $\mathcal{A}_{CP}^{dir}(B^0 \rightarrow \pi^0 \eta^{(\prime)})$ directly quoted from Ref. [8]

$$\mathcal{A}_{CP}^{dir}(B^0 \rightarrow \pi^0 \eta) = (-17.9_{-4.1}^{+5.2}{}_{-14.1}^{+7.9}{}_{-1.4}^{+1.2}{}_{-32.9}^{+33.4}) \times 10^{-2}, \quad (88)$$

$$\mathcal{A}_{CP}^{dir}(B^0 \rightarrow \pi^0 \eta') = (-19.2_{-4.3}^{+5.5}{}_{-7.8}^{+7.7}{}_{-3.3}^{+4.1}{}_{-35.8}^{+35.7}) \times 10^{-2}, \quad (89)$$

where the “default values” of the input parameters have been used [8], and the error sources are the same as those for the numerical results in Eqs. (77) and (78). Currently, no relevant experimental measurements for the CP-violating asymmetries of $B \rightarrow \pi^0 \eta^{(\prime)}$ decays are available. For the direct CP-violating asymmetries of $B^0 \rightarrow \pi^0 \eta^{(\prime)}$ decays, the theoretical predictions in pQCD and QCDF approach have the same sign, but the theoretical errors are clearly too large to make a meaningful comparison. One has to wait for the improvements in both the experimental measurements and the calculation of high order contributions.

If we integrate the time variable t , we will get the total CP asymmetry for $B^0 \rightarrow \pi^0 \eta^{(\prime)}$ decays,

$$A_{CP} = \frac{1}{1+x^2} A_{CP}^{dir} + \frac{x}{1+x^2} A_{CP}^{mix}, \quad (90)$$

where $x = \Delta m / \Gamma = 0.771$ for the $B^0 - \overline{B}^0$ mixing [28]. In Fig.8, we show the α -dependence of the total CP asymmetry A_{CP} for $B^0 \rightarrow \pi^0 \eta$ (solid curve) and $B^0 \rightarrow \pi^0 \eta'$ (solid curve)




FIG. 6: The direct CP asymmetry A_{CP}^{dir} (in percentage) of $B^0 \rightarrow \pi^0 \eta$ (solid curve) and $B^0 \rightarrow \pi^0 \eta'$ (dotted curve) as a function of CKM angle α .




FIG. 7: The mixing induced CP asymmetry A_{CP}^{mix} (in percentage) of $B^0 \rightarrow \pi^0 \eta$ (solid curve) and $B^0 \rightarrow \pi^0 \eta'$ (dotted curve) as a function of CKM angle α .

decay, respectively. Numerically, we found

$$\mathcal{A}_{CP}^{tot}(B^0 \rightarrow \pi^0 \eta) = (+3.7_{-7.3}^{+3.5}(\alpha)_{-3.9}^{+4.6}(\omega_b)_{-1.0}^{+0.3}(m_0^\pi)) \times 10^{-2}, \quad (91)$$

$$\mathcal{A}_{CP}^{tot}(B^0 \rightarrow \pi^0 \eta') = (+7.7_{-5.1}^{+1.9}(\alpha)_{-0.8}^{+0.3}(\omega_b)_{-5.1}^{+1.9}(m_0^\pi)) \times 10^{-2}. \quad (92)$$

D. Effects of possible gluonic component of η'

Up to now, we have not considered the possible contributions to the branching ratios and CP-violating asymmetries of $B \rightarrow \pi \eta'$ decays induced by the possible gluonic component of η' [29, 30, 31]. When $Z_{\eta'} \neq 0$, a decay amplitude $\mathcal{M}'$ will be produced by the gluonic component of η' . Such decay amplitude may construct or destruct with the ones




FIG. 8: The total CP asymmetry A_{cp}^{tot} (in percentage) of $B^0 \rightarrow \pi^0 \eta$ (solid curve) and $B^0 \rightarrow \pi^0 \eta'$ (dotted curve) as a function of CKM angle α .

from the $q\bar{q}$ ($q = u, d, s$) components of η' , the branching ratios of the decays in question may be increased or decreased accordingly.

Unfortunately, we currently do not know how to calculate this kind of contributions reliably. But we can treat it as an theoretical uncertainty. For $|M'/M(q\bar{q})| \sim 0.1 - 0.2$, for example, the resulted uncertainty for the branching ratios as given in Eqs.(66) and (68) will be around twenty to thirty percent.

From Eq. (2,66), one can see that the theoretical result of $Br(B^+ \rightarrow \pi^+ \eta')$ in the PQCD approach agree very well with the measured values within one standard deviation. Furthermore, the pQCD predictions for the branching ratios of $B \rightarrow \rho \eta^{(\prime)}$ decays also show very good agreement with the data [14]. We therefore believe that the gluonic admixture of η' should be small, and most possibly not as important as expected before.

As for the CP-violating asymmetries of $B \rightarrow \pi \eta'$ decays, the possible contributions of the gluonic components of the η' meson are largely cancelled in the ratio.

V. SUMMARY

In this paper, we calculate the branching ratios and CP-violating asymmetries of $B^0 \rightarrow \pi^0 \eta$, $B^0 \rightarrow \pi^0 \eta'$, $B^+ \rightarrow \pi^+ \eta$, and $B^+ \rightarrow \pi^+ \eta'$ decays in the PQCD factorization approach.

Besides the usual factorizable diagrams, the non-factorizable and annihilation diagrams are also calculated analytically. Although the non-factorizable and annihilation contributions are sub-leading for the branching ratios of the considered decays, but they are not negligible. Furthermore these diagrams provide the necessary strong phase required by a non-zero CP-violating asymmetry for the considered decays.

From our calculations and phenomenological analysis, we found the following results:

- From analytical calculations, the form factors for $B \rightarrow \eta$, $B \rightarrow \eta'$ and $B \rightarrow \pi$ transitions can be extracted. The PQCD results for these form factors are $F_{0,1}^{B \rightarrow \pi}(0) = 0.30$, $F_{0,1}^{B \rightarrow \eta}(0) = 0.15$ and $F_{0,1}^{B \rightarrow \eta'}(0) = 0.14$.

- For the CP-averaged branching ratios of the four considered decay modes, the theoretical predictions in PQCD approach are

$$\begin{aligned}
Br(B^+ \rightarrow \pi^+ \eta) &= (4.1_{-1.1}^{+1.5}) \times 10^{-6}, \\
Br(B^+ \rightarrow \pi^0 \eta^{(\prime)}) &= (2.4_{-0.6}^{+0.9}) \times 10^{-6}, \\
Br(B^0 \rightarrow \pi^0 \eta) &= (0.23 \pm 0.08) \times 10^{-6}, \\
Br(B^+ \rightarrow \pi^0 \eta^{(\prime)}) &= (0.19 \pm 0.05) \times 10^{-6},
\end{aligned} \tag{93}$$

where the various errors as specified in Eqs. (65-68) have been added in quadrature. Although the theoretical uncertainties are still large (can reach 40%), the leading PQCD predictions agree very well with the measured values or currently available experimental upper limits, and are also well consistent with the results obtained by employing the QCD factorization approach.

- For the CP-violating asymmetries, the theoretical predictions in PQCD approach are

$$\begin{aligned}
A_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta) &= (-37_{-7}^{+9}) \times 10^{-2}, \\
A_{CP}^{dir}(B^\pm \rightarrow \pi^\pm \eta') &= (-33_{-8}^{+7}) \times 10^{-2}, \\
A_{CP}^{dir}(B^0 \rightarrow \pi^0 \eta) &= (-42_{-13}^{+10}) \times 10^{-2}, \\
A_{CP}^{mix}(B^0 \rightarrow \pi^0 \eta) &= (+67_{-11}^{+5}) \times 10^{-2}, \\
A_{CP}^{dir}(B^0 \rightarrow \pi^0 \eta') &= (-36_{-10}^{+11}) \times 10^{-2}, \\
A_{CP}^{mix}(B^0 \rightarrow \pi^0 \eta') &= (+67_{-11}^{+5}) \times 10^{-2},
\end{aligned} \tag{94}$$

where the various errors as specified in Eqs. (75,76) and (84-87) have been added in quadrature. The uncertainties are around twenty to forty percent. And finally, the time integrated CP asymmetry for the neutral decay modes are

$$\begin{aligned}
\mathcal{A}_{CP}^{tot}(B^0 \rightarrow \pi^0 \eta) &= (+3.7_{-8.3}^{+5.8}) \times 10^{-2}, \\
\mathcal{A}_{CP}^{tot}(B^0 \rightarrow \pi^0 \eta') &= (+7.7_{-5.4}^{+2.7}) \times 10^{-2}.
\end{aligned} \tag{95}$$

- For $B^\pm \rightarrow \pi^\pm \eta$ decay, the pQCD prediction for $\mathcal{A}_{CP}^{dir}$ has the same sign with the measured value and consistent with the data within two standard deviations. For $B^\pm \rightarrow \pi^\pm \eta'$ decay, however, the pQCD prediction for $\mathcal{A}_{CP}^{dir}$ has an opposite sign with the measured value. Great improvements in both the theoretical calculations and experimental measurements are needed to clarify this discrepancy.
- From the very good agreement of the pQCD predictions for the CP-averaged branching ratios $Br(B^+ \rightarrow \pi^+ \eta^{(\prime)})$ and $Br(B^+ \rightarrow \rho^+ \eta^{(\prime)})$ [14] with the measured values, we believe that the gluonic admixture of η' should be small, and most possibly not as important as expected before.

Acknowledgments

We are very grateful to Li Ying for helpful discussions. This work is partly supported by the National Natural Science Foundation of China under Grant No.10275035, 10475085,

10575052, and by the Research Foundations of Jiangsu Education Committee and Nanjing Normal University under Grant No. 214080A916 and 2003102TSJB137.

APPENDIX A: RELATED FUNCTIONS

We show here the function h_i 's, coming from the Fourier transformations of $H^{(0)}$,

$$h_e(x_1, x_3, b_1, b_3) = K_0(\sqrt{x_1 x_3} m_B b_1) [\theta(b_1 - b_3) K_0(\sqrt{x_3} m_B b_1) I_0(\sqrt{x_3} m_B b_3) + \theta(b_3 - b_1) K_0(\sqrt{x_3} m_B b_3) I_0(\sqrt{x_3} m_B b_1)] S_t(x_3), \quad (\text{A1})$$

$$h_a(x_2, x_3, b_2, b_3) = K_0(i\sqrt{x_2 x_3} m_B b_3) [\theta(b_3 - b_2) K_0(i\sqrt{x_3} m_B b_3) I_0(i\sqrt{x_3} m_B b_2) + \theta(b_2 - b_3) K_0(i\sqrt{x_3} m_B b_2) I_0(i\sqrt{x_3} m_B b_3)] S_t(x_3), \quad (\text{A2})$$

$$h_f(x_1, x_2, x_3, b_1, b_2) = \left\{ \theta(b_2 - b_1) I_0(M_B \sqrt{x_1 x_3} b_1) K_0(M_B \sqrt{x_1 x_3} b_2) + (b_1 \leftrightarrow b_2) \right\} \cdot \left(\begin{array}{ll} K_0(M_B F_{(1)} b_1), & \text{for } F_{(1)}^2 > 0 \\ \frac{\pi i}{2} H_0^{(1)}(M_B \sqrt{|F_{(1)}^2|} b_1), & \text{for } F_{(1)}^2 < 0 \end{array} \right), \quad (\text{A3})$$

$$h_f^3(x_1, x_2, x_3, b_1, b_2) = \left\{ \theta(b_1 - b_2) K_0(i\sqrt{x_2 x_3} b_1 M_B) I_0(i\sqrt{x_2 x_3} b_2 M_B) + (b_1 \leftrightarrow b_2) \right\} \cdot \frac{\pi i}{2} H_0^{(1)}(\sqrt{x_1 + x_2 + x_3 - x_1 x_3 - x_2 x_3} b_1 M_B), \quad (\text{A4})$$

$$h_f^4(x_1, x_2, x_3, b_1, b_2) = \left\{ \theta(b_1 - b_2) K_0(i\sqrt{x_2 x_3} b_1 M_B) I_0(i\sqrt{x_2 x_3} b_2 M_B) + (b_1 \leftrightarrow b_2) \right\} \cdot \left(\begin{array}{ll} K_0(M_B F_{(2)} b_1), & \text{for } F_{(2)}^2 > 0 \\ \frac{\pi i}{2} H_0^{(1)}(M_B \sqrt{|F_{(2)}^2|} b_1), & \text{for } F_{(2)}^2 < 0 \end{array} \right), \quad (\text{A5})$$

where J_0 is the Bessel function and K_0 , I_0 are modified Bessel functions $K_0(-ix) = -(\pi/2)Y_0(x) + i(\pi/2)J_0(x)$, and $F_{(j)}$'s are defined by

$$F_{(1)}^2 = (x_1 - x_2)x_3, \quad (\text{A6})$$

$$F_{(2)}^2 = (x_1 - x_2)x_3. \quad (\text{A7})$$

The threshold resummation form factor $S_t(x_i)$ is adopted from Ref.[25]

$$S_t(x) = \frac{2^{1+2c} \Gamma(3/2 + c)}{\sqrt{\pi} \Gamma(1 + c)} [x(1 - x)]^c, \quad (\text{A8})$$

where the parameter $c = 0.3$. This function is normalized to unity.

The Sudakov factors used in the text are defined as

$$S_{ab}(t) = s\left(x_1 m_B/\sqrt{2}, b_1\right) + s\left(x_3 m_B/\sqrt{2}, b_3\right) + s\left((1-x_3)m_B/\sqrt{2}, b_3\right) - \frac{1}{\beta_1} \left[\ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_3\Lambda)} \right], \quad (\text{A9})$$

$$S_{cd}(t) = s\left(x_1 m_B/\sqrt{2}, b_1\right) + s\left(x_2 m_B/\sqrt{2}, b_2\right) + s\left((1-x_2)m_B/\sqrt{2}, b_2\right) + s\left(x_3 m_B/\sqrt{2}, b_1\right) + s\left((1-x_3)m_B/\sqrt{2}, b_1\right) - \frac{1}{\beta_1} \left[2 \ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_2\Lambda)} \right], \quad (\text{A10})$$

$$S_{ef}(t) = s\left(x_1 m_B/\sqrt{2}, b_1\right) + s\left(x_2 m_B/\sqrt{2}, b_2\right) + s\left((1-x_2)m_B/\sqrt{2}, b_2\right) + s\left(x_3 m_B/\sqrt{2}, b_2\right) + s\left((1-x_3)m_B/\sqrt{2}, b_2\right) - \frac{1}{\beta_1} \left[\ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + 2 \ln \frac{\ln(t/\Lambda)}{-\ln(b_2\Lambda)} \right], \quad (\text{A11})$$

$$S_{gh}(t) = s\left(x_2 m_B/\sqrt{2}, b_1\right) + s\left(x_3 m_B/\sqrt{2}, b_2\right) + s\left((1-x_2)m_B/\sqrt{2}, b_1\right) + s\left((1-x_3)m_B/\sqrt{2}, b_2\right) - \frac{1}{\beta_1} \left[\ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_2\Lambda)} \right], \quad (\text{A12})$$

where the function $s(q, b)$ are defined in the Appendix A of Ref.[9]. The scale t_i 's in the above equations are chosen as

$$t_e^1 = \max(\sqrt{x_3} m_B, 1/b_1, 1/b_3), \quad (\text{A13})$$

$$t_e^2 = \max(\sqrt{x_1} m_B, 1/b_1, 1/b_3), \quad (\text{A14})$$

$$t_e^3 = \max(\sqrt{x_3} m_B, 1/b_2, 1/b_3), \quad (\text{A15})$$

$$t_e^4 = \max(\sqrt{x_2} m_B, 1/b_2, 1/b_3), \quad (\text{A16})$$

$$t_f = \max(\sqrt{x_1 x_3} m_B, \sqrt{(x_1 - x_2)x_3} m_B, 1/b_1, 1/b_2), \quad (\text{A17})$$

$$t_f^3 = \max(\sqrt{x_1 + x_2 + x_3 - x_1 x_3 - x_2 x_3} m_B, \sqrt{x_2 x_3} m_B, 1/b_1, 1/b_2), \quad (\text{A18})$$

$$t_f^4 = \max(\sqrt{x_2 x_3} m_B, 1/b_1, 1/b_2). \quad (\text{A19})$$

-
- [1] I.I. Bigi and A.I. Sanda, *CP Violation* (Cambridge University Press, Cambridge, England, 2000); G.C. Branco, L. Lavoura and J.P. Silva, *CP Violation* (Oxford University Press, Oxford, England, 1999); R. Fleischer, Phys. Rep. **370** (2002) 537; T. Hurth, Rev. Mod. Phys. **75** (2003) 1159.
 - [2] M. Beneke, G. Buchalla, M. Neubert, and C.T. Sachrajda, Phys. Rev. Lett. **83** (1999) 914.
 - [3] C.-H. V. Chang and H.-n. Li, Phys. Rev. D **55** (1997) 5577; T.-W. Yeh and H.-n. Li, Phys. Rev. D **56** (1997) 1615.
 - [4] H.-n. Li, Prog. Part. & Nucl. Phys. **51** (2003) 85, and reference therein.
 - [5] G.P. Lepage and S. Brodsky, Phys. Rev. D **22** (1980) 2157; J. Botts and G. Sterman, Nucl. Phys. B **325** (1989) 62.
 - [6] D.S. Du, H.J. Gong, J.F. Sun, D.S. Yang, and G.H. Zhu, Phys. Rev. D **65** (2002) 074001, *ibid* **65** (2002) 094025; J.F. Sun, G.H. Zhu and D.S. Du, Phys. Rev. D **68** (2003) 054003.
 - [7] M.Z. Yang and Y.D. Yang, Nucl. Phys. B **609** (2001) 469; M. Beneke and M. Neubert, Nucl. Phys. B **651** (2003) 225.
 - [8] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003).
 - [9] C.-D. Lü, K. Ukai and M.Z. Yang, Phys. Rev. D **63** (2001) 074009.
 - [10] Y.-Y. Keum, H.-n. Li and A.I. Sanda, Phys. Lett. B **504** (2001) 6; Phys. Rev. D **63** (2001) 054008.
 - [11] H.-n. Li, Phys. Rev. D **64** (2001) 014019; S. Mishima, Phys. Lett. B **521** (2001) 252; C.-H. Chen, Y.-Y. Keum, and H.-n. Li, Phys. Rev. D **64** (2001) 112002; A.I. Sanda and K. Ukai, Prog. Theor. Phys. **107** (2002) 421; C.D. Lü, Eur. Phys. J. C **24** (2002) 121; C.-H. Chen, Y.-Y. Keum, and H.-n. Li, Phys. Rev. D **66** (2002) 054013; Y.-Y. Keum and A.I. Sanda, Phys. Rev. D **67** (2003) 054009.
 - [12] Y.-Y. Keum, T. Kurimoto, H.-n. Li, C.D. Lü, and A.I. Sanda, Phys. Rev. D **69** (2004) 094018; Y. Li and C.D. Lü, J. Phys. G **29** (2003) 2115; C.D. Lü, Phys. Rev. D **68** (2003) 097502; X.Q. Yu, Y. Li and C.D. Lü, Phys. Rev. D **71** (2005) 074026; C.D. Lü, Y.L. Shen and J. Zhu, Eur. Phys. J. C **41** (2005) 311; J. Zhu, Y.L. Shen and C.D. Lü, Phys. Rev. D **72** (2005) 054015; and hep-ph/0506316; Y. Li and C.D. Lü, Commun. Theor. Phys. **44** (2005) 659, hep-ph/0502038; and hep-ph/0508032; C.D. Lü, M. Matsumori, A.I. Sanda, and M.Z. Yang, Phys. Rev. D **72** (2005) 094005.
 - [13] Y. Li, C.D. Lü, Z.J. Xiao, and X.Q. Yu, Phys. Rev. D **70** (2004) 034009; Y. Li, C.D. Lü, and Z.J. Xiao, J. Phys. G **31** (2005) 273.
 - [14] X. Liu, H.S. Wang, Z.J. Xiao, L.B. Guo, and C.D. Lü, hep-ph/0509362.
 - [15] G. Buchalla, A.J. Buras, and M.E. Lautenbacher, Rev. Mod. Phys. **68** (1996) 1125.
 - [16] A. Ali, G. Kramer, and C.D. Lü, Phys. Rev. D **58** (1998) 094009.
 - [17] BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. D **70** (2004) 032006; BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **95** (2005) 131801.
 - [18] Belle Collaboration, P. Chang *et al.*, Phys. Rev. D **71** (2005) 091106(R); Belle Collaboration, K. Abe *et al.*, hep-ex/0508030 and hep-ex/0509016.
 - [19] Heavy Flavor Averaging Group, <http://www.slac.stanford.edu/xorg/hfag>.
 - [20] H.-n. Li, Phys. Rev. D **66** (2002) 094010.
 - [21] H.-n. Li and B. Tseng, Phys. Rev. D **57** (1998) 443.
 - [22] A.G. Grozin and M. Neubert, Phys. Rev. D **55** (1997) 272; M. Beneke and T. Feldmann,

- Nucl. Phys. B **592** (2001) 3.
- [23] M. Beneke, G. Buchalla, M. Neubert, and C.T. Sachrajda, Nucl. Phys. B **591** (2000) 313.
 - [24] C.D. Lü, M.Z. Yang, Eur. Phys. J. C **28** (2003) 515.
 - [25] T. Kurimoto, H.-n. Li, A.I. Sanda, Phys. Rev. D **65** (2002) 014007.
 - [26] C.-H. Chen and H.-n. Li, Phys. Rev. D **63** (2000) 014003.
 - [27] C.D. Lü, M.Z. Yang, Eur. Phys. J. C **23** (2002) 275.
 - [28] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592** (2004) 1.
 - [29] E. Kou, Phys. Rev. D **63** (2001) 054027.
 - [30] E. Kou and A.I. Sanda, Phys. Lett. B **525** (2002) 240.
 - [31] J.L. Rosner, Phys. Rev. D **27** (1983) 1101.
 - [32] V.M. Braun and I.E. Filyanov, Z. Phys. C **48** (1990) 239; P. Ball, J. High Energy Phys. 01 (1999) 010.
 - [33] P. Ball, J. High Energy Phys. 9809 (1998) 005; P. Ball, J. High Energy Phys. 9901 (1999) 010;
 - [34] T. Feldmann and P. Kroll, Eur. Phys. J. C **5** (1998) 327.
 - [35] V.M. Braun and I.E. Filyanov, Z. Phys. C **48** (1990) 239.
 - [36] P. Ball and V.M. Braun, Phys. Rev. D **58** (1998) 094016; P. Ball and R. Zwicky, Phys. Rev. D **71** (2005) 014015.
 - [37] J. Charles *et al.*, Eur. Phys. J. C **41** (2005) 1.
 - [38] A. Ali, G. Kramer and C.D. Lü, Phys. Rev. D **59** (1999) 014005.
 - [39] H.-n. Li, S. Mishima, A.I. Sanda, hep-ph/0508041.